

Updated butterflies of (Lepidoptera: Papilionoidia) Semuliki National Park following on from Davenport (1996) and Forbes (2018)

FAMILY PAPILIONIDAE Latreille, 1802	* New record (since Forbes, 2018) ^ Not recorded in DRC locations Ituri/Semliki Valley (Ducarme, 2025)
Subfamily Papilioninae Latreille, 1802	
Tribe Leptocircini Kirby, 1896	
Genus <i>Graphium</i> Scopoli, 1777	
<i>G.angolanus baronis</i> (Ungemach, 1932)	
<i>G.leonidas leonidas</i> (Fabricius, 1793)	
<i>G. polices polices</i> (Cramer, 1775)	
<i>G.antheus</i> (Cramer, 1779)	^
<i>G. ucalegon schoutedini</i> Berger, 1950	*
Tribe Papilionini Latreille, 1802	
Genus <i>Papilio</i> Linnaeus, 1758	
<i>P. mackinnoni mackinnoni</i> Sharpe, 1891	
<i>P. charopus montuosus</i> Joicey & Talbot, 1927	
<i>P. sosia pulchra</i> Berger, 1950	
<i>P. chrapkowskoides chrapkowskoides</i> Storace, 1952	
<i>P. nireus nireus</i> Linnaeus, 1758	
<i>P. cynorta</i> Fabricius, 1793	
<i>P. dardanus dardanus</i> Brown, 1776	
<i>P. phorcas congoanus</i> Rothschild, 1896	
<i>P. nobilis crippsianus</i> Stoneham, 1936	
<i>P. hesperus hesperus</i> Westwood, 1843	
<i>P. lormieri semlikana</i> Le Cerf, 1924	
<i>P. demodocus demodocus</i> Esper, 1798	
<i>P. mehowi mehowi</i> Dewitz, 1881	
<i>P. zenobia</i> Fabricius, 1775	
Total for the family Papilionidae	19
FAMILY HESPERIIDAE Latreille, 1809	
Subfamily Coeliadinae Evans, 1937	
Genus <i>Coeliades</i> Hübner, 1818	
<i>C. chalybe chalybe</i> (Westwood, 1852)	
<i>C. forestan forestan</i> (Stoll, 1782)	
<i>C. libeon</i> (Druce, 1875)	
Total for the subfamily Coeliadinae	3
Subfamily Pyrginae Mabille, 1878	
Tribe Tagiadini Mabille, 1878	
Genus <i>Eagris</i> Guenée, 1862	
<i>E.lucetia</i> (Hewitson, 1875)	

<i>E. tigris tigris</i> Evans, 1937	*
<i>E. nottoana nottoana</i> (Wallengren, 1857)	*
<i>E. decastigma purpura</i> Evans, 1937	
<i>E. tetrastigma tetrastigma</i> (Mabille, 1891)	*^
Genus <i>Calleagris</i> Aurivillius, 1925	
<i>C. hollandi</i> (Butler, 1897)	^
<i>C. lacteus lacteus</i> (Mabille, 1877)	
Genus <i>Tagiades</i> Hübner, 1819	
<i>T. flesus</i> (Fabricius, 1781)	
Genus <i>Abantis</i> Hopffer, 1855	
<i>A. iruma</i> Evans, 1951	*
Tribe Celaenorrhini Swinhoe, 1912	
Genus <i>Sarangesa</i> Moore, 1881	
<i>S. bouvieri</i> (Mabille, 1877)	*
<i>S. lucidella lucidella</i> (Mabille, 1891)	^
<i>S. brigida sanaga</i> Miller, 1964	
<i>S. tertullianus</i> (Fabricius, 1793)	*
<i>S. maculata</i> (Mabille, 1891)	*
<i>S. thecla mabira</i> Evans, 1956	*
<i>S. loelius</i> (Mabille, 1877)	*^
<i>S. haplopa</i> Swinhoe, 1907	*
Genus <i>Scopulifera</i> Moore, 1881	
<i>S. nigropunctata</i> (Bethune & Baker, 1908)	*
Genus <i>Bettonula</i> Libert & Larsen, 2014	
<i>B. bettoni bettoni</i> (Butler, 1902)	*
Genus <i>Apallaga</i> Strand, 1911	
<i>A. alluaudi</i> (Mabille & Boulet, 1916)	*
<i>A. fulgens montanus</i> Libert, 2014	*
<i>A. homeyeri fracta</i> Libert, 2014	*
<i>A. darlustris</i> Libert, 2014	*
<i>A. inexpectata</i> Libert, 2014	*
<i>A. boadicea howarthi</i> (Berger, 1976)	*
Genus <i>Eretis</i> Mabille, 1891	
<i>E. buamba</i> Evans, 1937	*
<i>E. melania</i> Mabille, 1891	*
<i>E. artorius</i> Larsen & Cock, 2011	*
<i>E. cameroni</i> Evans, 1937	*
Genus <i>Celaenorrhinus</i> Hübner, 1819	
<i>C. proxima proxima</i> (Mabille, 1877)	*
<i>C. perimitans</i> Libert, 2014	*
<i>C. macrostictus</i> (Holland, 1893)	*
Tribe Carcharodini Verity, 1940	

Genus <i>Spialia</i> Swinhoe, 1912	
<i>S. ploetzi ploetzi</i> (Aurivillius, 1891)	*
<i>S. spio</i> (Linnaeus, 1764)	*
Total for the subfamily Pyrginae	34
Subfamily Hesperinae Latreille, 1809	
Tribe Aeromachini Tutt, 1906	
Genus <i>Lepella</i> Evans, 1937	
<i>L. lepeletier</i> (Latreille, 1824)	*
Genus <i>Prosopalpus</i> Holland, 1896	
<i>P. styła</i> Evans, 1937	*
Tribe Astictopterini Swinhoe, 1912	
Genus <i>Kedestes</i> Watson, 1893	
<i>K. rogersi</i> Druce, 1907	^
Genus <i>Gorgyra</i> Holland, 1896	
<i>G. mocquersii</i> Holland, 1896	*
<i>G. diversata</i> Evans, 1937	*
Genus <i>Teniorhinus</i> Holland, 1892	
<i>T. ignita</i> (Mabille, 1877)	
<i>T. niger</i> (Druce, 1910)	*
Genus <i>Xanthodisca</i> Aurivillius, 1925	
<i>X. vibius</i> (Hewitson, 1878)	*
<i>X. astrape</i> (Holland, 1892)	*
Genus <i>Hypoleucis</i> Mabille, 1891	
<i>H. tripunctata draga</i> Evans, 1937	
<i>H. dacena</i> Hewitson, 1876	*
<i>H. ophiusa ophir</i> Evans, 1937	*
Genus <i>Andronymus</i> Holland, 1896	
<i>A. neander neander</i> Plötz, 1884	
<i>A. gander</i> Evans, 1947	*
<i>A. caesar philander</i> Hopffer, 1855	*
Genus <i>Zophopetes</i> Mabille, 1904	
<i>Z. dysmephila</i> (Trimen, 1868)	
<i>Z. cerymica</i> Hewitson, 1867	*
<i>Z. ganda</i> Evans, 1937	*
<i>Z. nobilor</i> Holland, 1896	^
Genus <i>Fresna</i> Evans, 1937	
<i>F. nyassae</i> Hewitson, 1878	*
<i>F. netopha</i> (Hewitson, 1878)	*
Tribe Baorini Doherty, 1886	
Genus <i>Zenonia</i> Evans, 1935	
<i>Z. zeno</i> (Trimen, 1864)	*
<i>Z. crasta</i> Evans, 1937	*

Genus <i>Pelopidas</i> Walker, 1870	
<i>P. thrax</i> (Hübner, 1821)	*
<i>P. mathias mathias</i> (Fabricius, 1798)	*
Genus <i>Borbo</i> Evans, 1949	
<i>B. lugens</i> (Hopffer, 1855)	*
<i>B. detecta</i> (Trimen, 1893)	*
<i>B. borbonica borbonica</i> (Boisduval, 1833)	*
<i>B. fatuellus fatuellus</i> (Hopffer, 1855)	*
<i>B. fallax</i> Gaede, 1916	
<i>B. micans</i> (Holland, 1896)	*
Tribe Ceratrichiini Grishin, 2019	
Genus <i>Ceratrichia</i> Butler, 1870	
<i>C. semlikensis</i> Joicey & Talbot, 1921	
<i>C. clara medea</i> Evans, 1937	^
<i>C. hollandi</i> Bethune-Baker, 1908	*
<i>C. wollastoni</i> Heron, 1909	
<i>C. enta</i> Evans, 1947	*
Genus <i>Pardaleodes</i> Butler, 1870	
<i>P. tibullus torensis</i> Bethune-Baker, 1906	
<i>P. sator pusiella</i> Mabille, 1877	
<i>P. bule</i> Holland, 1896	*
<i>P. incerta incerta</i> (Snellen, 1872)	
Genus <i>Meza</i> Hemming, 1939	
<i>M. cybeutes pallida</i> (Evans, 1937)	
Genus <i>Ankola</i> Evans, 1937	
<i>A. fan</i> (Holland, 1894)	*
Tribe Astictopterini Swinhoe, 1912	
Genus <i>Ceratricula</i> Larsen, 2013	
<i>C. semilutea congdoni</i> Larsen, 2013	
Genus <i>Osmodes</i> Holland, 1892	
<i>O. laronia</i> (Hewitson, 1868)	
<i>O. thora</i> (Plötz, 1884)	
<i>O. banghaasi</i> Holland, 1896	*^
<i>O. adonia</i> Evans, 1937	*
<i>O. adosus</i> (Mabille, 1889)	*
<i>O. lux</i> Holland, 1892	*
<i>O. hollandi</i> Evans, 1937	*
Genus <i>Gamia</i> Holland, 1896	
<i>G. shelleyi</i> (Sharpe, 1890)	
Genus <i>Caenides</i> Holland, 1896	
<i>C. dacela</i> (Hewitson, 1876)	
<i>C. kangvensis</i> Holland, 1896	*

Genus <i>Monza</i> Evans, 1937	
<i>M. punctata crola</i> Evans, 1937	
<i>M. cretacea</i> (Snellen, 1872)	*
<i>M. alberti</i> (Holland, 1896)	*
Genus <i>Rhabdomantis</i> Holland, 1896	
<i>R. galatia</i> (Hewitson, 1868)	*
Genus <i>Acleros</i> Mabille, 1885	
<i>A. neavei</i> Evans, 1937	^
<i>A. ploetzi</i> Mabille, 1889	
<i>A. nigrapex</i> Strand, 1913	*
<i>A. bibundica</i> Strand, 1913	*
<i>A. olaus</i> Plötz, 1884	*
Genus <i>Semalea</i> Holland, 1896	
<i>S. pulvina</i> (Plötz, 1879)	*
<i>S. sextilis</i> (Plötz, 1886)	*
Genus <i>Paronymus</i> Aurivillius, 1925	
<i>P. ligora</i> (Hewitson, 1876)	*
Genus <i>Chondrolepis</i> Mabille, 1904	
<i>C. niveicornis</i> (Plötz, 1883)	*
Genus <i>Mopala</i> Evans, 1937	
<i>M. orma</i> (Plötz, 1879)	*
Genus <i>Pteroteinon</i> Watson, 1893	
<i>P. caenira</i> (Hewitson, 1867)	*
<i>P. capronnieri</i> (Plötz, 1879)	*
Tribe Gretnini Grisham, 2019	
Genus <i>Gretna</i> Evans, 1937	
<i>G. cylinda</i> (Hewitson, 1876)	
<i>G. zarembo jacksoni</i> Evans, 1937	*^
<i>G. carmen carmen</i> Evans, 1937	*
Total for the subfamily Hesperinae	72
Total for the family Hesperidae	109
FAMILY NYMPHALIDAE Rafinesque, 1815	
Subfamily Danainae Boisduval, 1833	
Tribe Danaini Boisduval, 1833	
Genus <i>Danaus</i> Kluk, 1780	
<i>D. chrysippus chrysippus</i> (Linnaeus, 1758)	
Genus <i>Tirumala</i> Moore, 1880	
<i>T. petiverana</i> (Doubleday, 1847)	
<i>T. formosa mercedonia</i> (Karsch, 1894)	
Genus <i>Amauris</i> Hübner, 1816	
<i>A. niavius niavius</i> (Linnaeus, 1758)	
<i>A. tartarea tartarea</i> (Mabille, 1876)	

<i>A. hecate hecate</i> (Butler, 1866)	
<i>A. albimaculata magnimacula</i> Rebel, 1914	
<i>A. crawshayi oscarus</i> Thurau, 1904	
Total for the subfamily Danainae	8
Subfamily Libytheinae Boisduval, 1833	
<i>L. labdaca</i> Westwood, 1851	
Total for the subfamily Libytheinae	1
Subfamily Satyrinae Boisduval, 1833	
Tribe Elymniini Herrich-Schaeffer, 1864	
Genus <i>Elymnias</i> Hübner, 1816	
<i>E. bammakoo ratrayi</i> Sharpe, 1902	
Tribe Melanitini Reuter, 1896	
Genus <i>Melanitis</i> Fabricius, 1807	
<i>M. leda</i> (Linnaeus, 1758)	
<i>M. libya</i> Distant, 1882	
Genus <i>Ducarmeia</i> Pyrcz, 2020	
<i>D. ansorgei</i> Rothschild, 1904	
Genus <i>Gnophodes</i> Doubleday, 1849	
<i>G. parmeno</i> Doubleday, 1849	
Genus <i>Haydonia</i> Pyrcz & Collins, 2020	
<i>H. chelys</i> (Fabricius, 1793)	
Tribe Satyrini Boisduval, 1833	
Genus <i>Bicyclus</i> Kirby, 1871	
<i>B. alboplaga</i> (Rebel, 1914)	+
<i>B. angulosa angulosa</i> (Butler, 1868)	
<i>B. buea</i> (Strand, 1912)	^
<i>B. campina campina</i> (Aurivillius, 1901)	^
<i>B. hyperanthus</i> (Bethune-Baker, 1908)	^
<i>B. graueri graueri</i> (Rebel, 1914)	
<i>B. sebetus</i> (Hewitson, 1877)	
<i>B. mesogena ugandae</i> (Riley, 1926)	
<i>B. mesogenina</i> Grünberg, (1911)	*^
<i>B. sambulos sambulos</i> Hewitson, 1877	
<i>B. collinsi</i> Aduse-Poku, Lees & Wahlberg, 2016	
<i>B. auricruda fulgida</i> Fox, 1963	
<i>B. vulgaris</i> (Butler, 1868)	
<i>B. jefferyi</i> Fox, 1963	
<i>B. moyses</i> Condamin and Fox, 1964	*
<i>B. sandace</i> (Hewitson, 1877)	
<i>B. sanaos</i> (Hewitson, 1866)	
<i>B. sophrosyne sophrosyne</i> (Plötz, 1880)	
<i>B. mollitia</i> (Karsch, 1895)	

<i>B. smithi smithi</i> (Aurivillius, 1899)	
<i>B. golo</i> (Aurivillius, 1893)	
<i>B. ignobilis</i> (Butler, 1870)	
<i>B. dentata</i> Sharpe, 1898	*
<i>B. safitza safitza</i> (Hewitson, 1850)	
<i>B. campus</i> (Karsch, 1893)	
<i>B. funebris</i> (Guérin-Méneville, 1844)	
<i>B. procora</i> (Karsch, 1893)	
<i>B. uniformis</i> (Bethune-Baker, 1908)	
Genus <i>Brakefieldia</i> Aduse-Poku, Lees & Wahlberg, 2016	
<i>B. phaea phaea</i> (Karsch, 1894)	
<i>B. peitho reducta</i> (Libert, 2006)	
Genus <i>Ypthima</i> Hübner, 1818	
<i>Y. asterope asterope</i> (Klug, 1832)	
<i>Y. doleta</i> Kirby, 1880	
<i>Y. albida albida</i> Butler, 1888	
Genus <i>Ypthimomorpha</i> van Son, 1955	
<i>Y. itonia</i> (Hewitson, 1865)	
Total for the subfamily Satyrinae	40
Subfamily Charaxinae Guenée, 1865	
Tribe Pallini Rydon, 1971	
Genus <i>Palla</i> Hübner, 1819	
<i>P. violinitens bwamba</i> van Someren, 1975	
Tribe Charaxini Guenée, 1865	
Genus <i>Charaxes</i> Oechsenheimer, 1816	
<i>C. varanes vologeses</i> (Mabille, 1876)	
<i>C. fulvescens monitor</i> Rothschild, 1900	
<i>C. acuminatus kigezia</i> van Someren, 1963	
<i>C. candiope</i> (Godart, 1824)	
<i>C. protoclea nothodes</i> Jordan, 1911	
<i>C. cynthia kinduana</i> Le Cerf, 1923	
<i>C. lucretius maximus</i> van Someren, 1971	
<i>C. epijasius</i> Reiche, 1850	
<i>C. castor castor</i> (Cramer, 1775)	
<i>C. brutus angustus</i> Rothschild, 1900	
<i>C. pollux pollux</i> (Cramer, 1775)	
<i>C. eudoxus mechowii</i> Rothschild, 1900	
<i>C. numenes aequatorialis</i> van Someren, 1972	
<i>C. tiridates tiridatinus</i> Röber, 1936	
<i>C. bipunctatus ugandensis</i> van Someren, 1972	
<i>C. smaragdalis smaragdalis</i> Butler, 1866	
<i>C. imperialis ugandicus</i> van Someren, 1972	

<i>C. ameliae victoriae</i> van Someren, 1972	
<i>C. pythodoris pythodoris</i> Hewitson, 1873	
<i>C. hadrianus</i> Ward, 1871	
<i>C. zingha</i> (Stoll, 1780)	
<i>C. etesipe etesipe</i> (Godart, 1824)	
<i>C. anticlea adusta</i> Rothschild, 1900	
<i>C. baumanni bwamba</i> van Someren, 1971	
<i>C. hildebrandti katangensis</i> Talbot, 1928	
<i>C. virilis lenis</i> Henning, 1989	
<i>C. catachrous</i> van Someren & Jackson, 1952	
<i>C. etheocles carpenteri</i> van Someren & Jackson, 1957	
<i>C. cedreatis</i> Hewitson, 1874	
<i>C. viola picta</i> van Someren & Jackson, 1952	
<i>C. pleione bebra</i> Rothschild, 1900	
<i>C. paphianus subpallida</i> Joicey & Talbot, 1925	
<i>C. kahlden</i> Homeyer & Dewitz, 1882	
<i>C. nichetes nichetes</i> Grose-Smith, 1883	
<i>C. lycurgus</i> (Fabricius, 1793)	
<i>C. zelica depuncta</i> Joicey & Talbot, 1921	
<i>C. trajanus vansomeri</i> (Poulton, 1929)	
<i>C. eurinome lequeuxi</i> (Turlin, 2009)	
<i>C. crossleyi magnifica</i> (Rebel, 1914)	
<i>C. eupale latimargo</i> Joicey & Talbot, 1921	
<i>C. minor</i> Joicey & Talbot, 1921	
<i>C. subornatus subornatus</i> Schultze, 1916	*^
Total for the subfamily Charaxinae	43
Subfamily Nymphalinae Rafinesque, 1815	
Tribe Kallimini Doherty, 1886	
Genus <i>Catacroptera</i> Karsch, 1894	
<i>C. cloanthe cloanthe</i> (Stoll, 1781)	
Tribe Junoniini Reuter, 1896	
Genus <i>Junonia</i> Hübner, 1819	
<i>J. chorimene</i> (Guérin-Ménéville, 1844)	
<i>J. gregorii</i> Butler, 1896	
<i>J. terea tereoides</i> (Butler, 1901)	
<i>J. sophia infracta</i> (Butler, 1888)	
<i>J. westermanni westermanni</i> (Westwood, 1870)	
<i>J. oenone oenone</i> (Linnaeus, 1758)	
<i>J. stygia</i> (Aurivillius, 1894)	
<i>J. ansorgei</i> (Rothschild, 1899)	
Genus <i>Salamis</i> Boisduval, 1833	
<i>S. cacta cacta</i> (Fabricius, 1793)	

Genus <i>Protoponiomorpha</i> Wallengren, 1857	
<i>P. temora temora</i> (Felder & Felder, 1867)	
<i>P. parhassus</i> (Drury, 1782)	
<i>P. anacardii nebulosa</i> (Trimen, 1881)	
Genus <i>Precis</i> Hübner, 1819	
<i>P. pelarga</i> (Fabricius, 1775)	
<i>P. actia</i> Distant, 1880	
<i>P. antilope</i> (Feisthamel, 1850)	^
<i>P. tugela pyriformis</i> (Butler, 1896)	
Genus <i>Hypolimnas</i> Hübner, 1819	
<i>H. misippus</i> (Linnaeus, 1764)	
<i>H. salmaceis salmaceis</i> (Drury, 1773)	
<i>H. monteironis monteironis</i> Druce, 1874	
<i>H. anthedon anthedon</i> (Doubleday, 1845)	
Tribe <i>incertae sedis</i>	
Genus <i>Kallimoides</i> Shirôzu & Nakanishi, 1984	
<i>K. rumia ratrayi</i> (Sharpe, 1904)	
Genus <i>Vanessula</i> Dewitz, 1887	
<i>V. milca latifascia</i> Joicey & Talbot, 1928	
Total for the subfamily Nymphalinae	23
Subfamily Cyrestinae Guenée, 1865	
Genus <i>Cyrestis</i> Boisduval, 1832	
<i>C. camillus camillus</i> (Fabricius, 1781)	
Total for the subfamily Cyrestinae	1
Subfamily Biblidinae Boisduval, 1833	
Tribe Biblidini Boisduval, 1833	
Genus <i>Byblia</i> Hübner, 1819	
<i>B. anvata craueri</i> Aurivillius, 1894	
<i>B. ilithyia</i> (Drury, 1773)	^
Genus <i>Mesoxantha</i> Aurivillius, 1899	
<i>M. ethosea ethesoides</i> Rebel, 1914	
Genus <i>Ariadne</i> Horsfield, 1829	
<i>A. enotrea suffusa</i> (Joicey & Talbot, 1921)	
<i>A. pagenstecheri</i> (Suffert, 1904)	^
<i>A. albifascia</i> (Joicey & Talbot, 1921)	*
Genus <i>Neptidopsis</i> Aurivillius, 1899	
<i>N. ophione nucleata</i> Grünberg, 1911	
Genus <i>Eurytela</i> Boisduval, 1833	
<i>E. hiarbas hiarbas</i> (Drury, 1782)	
<i>E. dryope angulata</i> Aurivillius, 1899	
Tribe Epicaliini Guenée, 1865	
Genus <i>Sevenia</i> Koçak, 1996	

<i>S. occidentaliu occidentaliu</i> (Mabille, 1876)	
<i>S. garega</i> (Karsch, 1892)	
<i>S. boisduvali omissa</i> (Rothschild, 1918)	
<i>S. umbrina</i> (Karsch, 1892)	*^
Total for the subfamily Biblidinae	13
Subfamily Apaturinae Boisduval, 1840	
Genus <i>Apaturopsis</i> Aurivillius, 1899	
<i>A. cleochares cleochares</i> (Hewitson, 1873)	
Total for the subfamily Apaturinae	1
Subfamily Heliconiinae Swainson, 1822	
Tribe Acraeini Boisduval, 1833	
Subgenus <i>Acraea</i>	
Genus <i>Acraea</i> Fabricius, 1807	
<i>A. kraka</i> Aurivillius, 1893	
<i>A. rogersi</i> Hewitson, 1873	
<i>A. egina egina</i> (Cramer, 1775)	
<i>A. abdera abdera</i> Hewitson, 1852	
<i>A. cepheus</i> (Linnaeus, 1758)	
<i>A. asboloplintha asboloplintha</i> Karsch, 1894	
<i>A. zetes zetes</i> (Linnaeus, 1758)	
<i>A. leucographa leucographa</i> Ribbe, 1889	
<i>A. quirina quirina</i> (Fabricius, 1781)	
<i>A. endoscuta</i> (Le Doux, 1928)	*
<i>A. eltringhami</i> (Joicey and Talbot, 1921)	*
<i>A. epaea epaea</i> (Cramer, 1779)	
<i>A. paragea</i> Grose-Smith, 1900	
<i>A. schubotzi</i> (Grünberg, 1911)	
<i>A. ducarme</i> Bernaud & Pierre, 2012	
<i>A. poggei nelsoni</i> Grose-Smith & Kirby, 1892	
<i>A. epiprotea</i> (Butler, 1874)	
<i>A. alcinoe alcinoe</i> (Felder & Felder, 1865)	
<i>A. consanguinea consanguinea</i> (Aurivillius, 1893)	
<i>A. pseuderyta</i> Godman & Salvin, 1890	
<i>A. macarista macarista</i> (Sharpe, 1906)	
<i>A. macaria hemileuca</i> (Jordan, 1914)	
Subgenus <i>Actinote</i>	
<i>A. perenna perenna</i> (Doubleday, 1847)	
<i>A. cinerea</i> (Neave, 1904)	
<i>A. orinata</i> (Oberthür, 1893)	
<i>A. penelope penelope</i> (Staudinger, 1896)	
<i>A. quirinalis</i> (Grose Smith, 1900)	
<i>A. parrhasia servona</i> (Godart, 1819)	

<i>A. peneleos pelasgia</i> (Grose-Smith, 1900)	
<i>A. grosvenori</i> (Eltringham, 1912)	
<i>A. oreas oreas</i> (Sharpe, 1891)	
<i>A. iturina</i> (Grose-Smith, 1890)	
<i>A. humilis</i> (Sharpe, 1897)	*
<i>A. pharsalus pharsalus</i> (Ward, 1871)	
<i>A. encendon encendon</i> (Linnaeus, 1758)	
<i>A. encedana</i> (Pierre, 1976)	
<i>A. jodutta jodutta</i> (Fabricius, 1793)	
<i>A. aurivillii aurivillii</i> (Staudinger, 1896)	
<i>A. alciopoides</i> (Joicey & Talbot, 1921)	
<i>A. lycoa</i> (Godart, 1819)	
<i>A. serena</i> (Fabricius, 1775)	
<i>A. acerata</i> (Hewitson, 1874)	
<i>A. ventura</i> (Hewitson, 1877)	*
<i>A. althoffi rubrofasciata</i> (Aurivillius, 1895)	
<i>A. uvui uvui</i> (Grose-Smith, 1890)	
<i>A. bonasia bonasia</i> (Fabricius, 1775)	
<i>A. cabira</i> (Hopffer, 1855)	*
<i>A. sotikensis sotikensis</i> (Sharpe, 1892)	*
<i>A. viviana</i> (Staudinger, 1896)	
<i>A. orestia orestia</i> (Hewitson, 1874)	
Tribe Vagrantini Pinratana & Eliot, 1996	
Genus <i>Lachnoptera</i> Doubleday, 1847	
<i>L. anticlia</i> (Hübner, 1819)	
Genus <i>Phalanta</i> Horsfield, 1829	
<i>P. phalantha aethiopica</i> (Rothschild & Jordan, 1903)	
<i>P. eurytis eurytis</i> (Doubleday, 1847)	
Tribe Argynnini Swainson, 1833	
Genus <i>Issoria</i> Hübner, 1819	
<i>I. baumanni excelsior</i> (Butler, 1896)	
Total for the subfamily Heliconiinae	54
Subfamily Limenitidinae Behr, 1864	
Tribe Pseudoneptini Dhungel & Wahlberg, 2018	
Genus <i>Pseudoneptis</i> Snellen, 1882	
<i>P. bugandensis ianthe</i> Hemming, 1964	
Tribe Cymothoini Dhungel & Wahlberg, 2018	
Genus <i>Cymothoe</i> Hübner, 1819	
<i>C. beckeri theodosia</i> Staudinger, 1890	
<i>C. confusa</i> Aurivillius, 1887	
<i>C. cyclades</i> (Ward, 1871)	
<i>C. ochreata</i> Grose Smith, 1890	

<i>C. lurida tristis</i> Overlaet, 1952	
<i>C. colmanti</i> Aurivillius, 1898	
<i>C. haynae diphya</i> Karsch, 1894	
<i>C. herminia johnstoni</i> (Butler, 1902)	
<i>C. caenis</i> (Drury, 1773)	
<i>C. coccinata vrydaghi</i> Overlaet, 1944	
<i>C. reginaeelisabethae reginaeelisabethae</i> Holland, 1920	
<i>C. distincta kivuensis</i> Overlaet, 1952	*
<i>C. sangaris sangaris</i> (Godart, 1824)	
<i>C. hobarti mwamikazi</i> Overlaet, 1952	
<i>C. jodutta mostinckxi</i> Overlaet, 1952	
<i>C. meridionalis ghesquierei</i> Overlaet, 1944	
Genus <i>Harma</i> Doubleday, 1848	
<i>H. theobene superna</i> (Fox, 1968)	
Tribe Neptini Newman, 1870	
Genus <i>Neptis</i> Fabricius, 1807	
<i>N. saclava marpessa</i> Hopffer, 1855	
<i>N. conspicua</i> Neave, 1904	
<i>N. constantiae constantiae</i> Carcasson, 1961	
<i>N. metanira</i> Holland, 1892	
<i>N. metella metella</i> (Doubleday, 1850)	
<i>N. nysiades</i> Hewitson, 1868	
<i>N. nemetes margueriteae</i> Fox, 1968	
<i>N. nicomedes</i> Hewitson, 1874	
<i>N. strigata kakamega</i> Collins & Larsen, 1996	
<i>N. nicoteles</i> Hewitson, 1874	
<i>N. melicerta parallela</i> Collins & Larsen, 1996	
<i>N. lermanni</i> Aurivillius, 1896	
<i>N. trigonophora intermedia</i> Schultze, 1920	
<i>N. kiriakoffi</i> Overlaet, 1955	^
<i>N. serena serena</i> Overlaet, 1955	
<i>N. morosa</i> Overlaet, 1955	
<i>N. seeldrayersi</i> Aurivillius, 1895	
<i>N. alta</i> Overlaet, 1955	
<i>N. ochracea ochracea</i> Neave, 1904	
<i>N. continuata</i> Holland, 1892	
Tribe Adoliadini Doubleday, 1845	
Genus <i>Evena</i> Westwood, 1850	
<i>E. crithea</i> (Drury, 1773)	
<i>E. angustatum</i> (Felder & Felder, 1867)	
<i>E. oberthueri</i> (Karsch, 1894)	
Genus <i>Aterica</i> Boisduval, 1833	

<i>A. galene extensa</i> Heron, 1909	
Genus <i>Pseudargynnis</i> Karsch, 1892	
<i>P. hegemone</i> (Godart, 1819)	
Genus <i>Cynandra</i> Schatz, 1887	
<i>C. opis bernardii</i> Lagnel, 1967	
Genus <i>Euryphura</i> Staudinger, 1891	
<i>E. chalcis chalcis</i> (Felder & Felder, 1860)	
<i>E. plautilla</i> (Hewitson, 1865)	
Genus <i>Euphaedra</i> Hübner, 1819	
<i>E. medon fraudata</i> Thureau, 1904	
<i>E. zaddachi zaddachii</i> Dewitz, 1879	
<i>E. mirabilis nubila</i> Hecq, 1986	
<i>E. caerulescens caerulescens</i> Grose Smith, 1890	
<i>E. ansorgei</i> Hecq, 1974	*
<i>E. christyi</i> Sharpe, 1904	^
<i>E. eberti hamus</i> Berger, 1940	
<i>E. ueleana</i> Hecq, 1982	*
<i>E. diffusa diffusa</i> Gaede, 1916	
<i>E. imitans</i> Holland, 1893	
<i>E. sarita sarita</i> (Sharpe, 1891)	
<i>E. sinuosa sinuosa</i> Hecq, 1974	*
<i>E. viridicaerulea inanoides</i> Holland, 1920	
<i>E. intermedia</i> Rebel, 1914	*
<i>E. preussi preussi</i> Staudinger, 1891	
<i>E. procera</i> Hecq, 1984	*
<i>E. vicina viscina</i> Hecq, 1984	*
<i>E. albofasciata</i> Rebel, 1914	
<i>E. eleus eleus</i> (Drury, 1782)	
<i>E. simplex</i> Hecq, 1978	*
<i>E. hybrida</i> Hecq, 1978	
<i>E. alacris</i> Hecq, 1978	
<i>E. rattrayi rattrayi</i> Sharpe, 1904	
<i>E. alternus</i> van Someren, 1935	*^
<i>E. ruspina</i> (Hewitson, 1865)	
<i>E. edwardsii edwardsi</i> van der Hoeven, 1845	
<i>E. harpalyce spatiosa</i> (Mabille, 1876)	
Genus <i>Euriphene</i> Boisduval, 1847	
<i>E. ribensis</i> (Ward, 1871)	
<i>E. atossa australis</i> d'Abrera, 1980	
<i>E. amaranta</i> (Karsch, 1894)	
<i>E. saphirina saphirina</i> (Karsch, 1894)	
<i>E. doriclea ducarme</i> Hecq, 1994	*

Genus <i>Bebearia</i> Hemming, 1960	
<i>B. carshena</i> (Hewitson, 1871)	
<i>B. absolon absolon</i> (Lathy, 1906)	
<i>B. abesa pandera</i> Hecq, 1987	
<i>B. zonara</i> (Butler, 1871)	
<i>B. mandinga beni</i> Hecq, 1990	
<i>B. oxione squalida</i> (Talbot, 1928)	
<i>B. partita</i> (Aurivillius, 1895)	
<i>B. barce maculata</i> (Aurivillius, 1912)	
<i>B. plistonax</i> (Hewitson, 1874)	
<i>B. cocalia badiana</i> (Rebel, 1914): <i>B. cocalia katera</i> van Someren, 1939: <i>B. cocalia continentalis</i> (Hecq, 1988)	
<i>B. sophus aruunda</i> Overlaet, 1955	
<i>B. seeldrayersi</i> (Aurivillius, 1899)	
<i>B. brunhilda iturina</i> (Karsch, 1894)	
<i>B. dowsetti</i> (Hecq, 1990)	*
<i>B. laetitioides</i> (Joicey & Talbot, 1921)	
<i>B. congolensis</i> (Capronnier, 1889)	*
<i>B. flaminia flaminia</i> (Staudinger, 1891)	
<i>B. chloeropsis</i> (Bethune-Baker 1908)	*
Genus <i>Hamanumida</i> Hübner, 1819	
<i>H. daedalus</i> (Fabricius, 1775)	
Tribe Pseudacraeini Dhungel & Wahlberg, 2018	
Genus <i>Pseudacraea</i> Westwood, 1850	
<i>P. eurytus eurytus</i> (Linnaeus, 1758)	
<i>P. lucretia protracta</i> (Butler, 1875)	
<i>P. semire</i> (Cramer, 1779)	
<i>P. warburgi</i> Aurivillius, 1892	
Total for the subfamily Limenitidinae	101
Total for the family Nymphalidae	285
FAMILY PIERIDAE Swainson, 1820	
Subfamily Pseudopontiinae Reuter, 1896	
Genus <i>Pseudopontia</i> Plötz, 1870	
<i>P. mabira</i> Mitter & Collins, 2011	
Total for subfamily Pseudopontiinae	1
Subfamily Coliadinae Swainson, 1821	
Subgenus <i>Eurema</i>	
Genus <i>Eurema</i> Hübner, 1819	
<i>E. hecabe solifera</i> (Butler, 1875)	
<i>E. floricola leonis</i> (Butler, 1886)	
<i>E. senegalensis</i> (Boisduval, 1836)	
<i>E. desjardinsi regularis</i> (Butler, 1876)	

<i>E. hapale</i> (Mabille, 1882)	
<i>E. brigitta brigitta</i> (Stoll, 1790)	*
<i>E. mandarinula</i> (Holland, 1892)	*
Genus <i>Catopsilia</i> Hübner, 1819	
<i>C. florella</i> (Fabricius, 1775)	
Total for the subfamily Coliadinae	8
Subfamily Pierinae Swainson, 1820	
Tribe Leptosiaini Braby, 2014	
Genus <i>Leptosia</i> Hübner, 1818	
<i>L. wigginsi wigginsi</i> (Dixey, 1915)	
<i>L. nupta pseudonupta</i> Bernardi, 1959	
<i>L. alcesta inalcesta</i> Bernardi, 1959	
<i>L. hybrida somereni</i> Bernardi, 1959	*
Tribe Nephroniini Braby, 2014	
Genus <i>Nepheronia</i> Butler, 1870	
<i>N. pharis silvanus</i> (Stoneham, 1957)	
<i>N. thalassina verulanus</i> (Ward, 1871)	
<i>N. argia argia</i> (Fabricius, 1775)	
<i>N. buquetii buquetii</i> (Boisduval, 1836)	
Tribe Teracolini Reuter, 1896	
Genus <i>Colotis</i> Hübner, 1819	
<i>C. euipe complexivus</i> (Butler, 1886)	
<i>C. hetaera ankolensis</i> Stoneham, 1940	
<i>C. elgonensis basilewskyi</i> Berger, 1956	
<i>C. antevippe zera</i> (Lucas, 1852)	
<i>C. aurigineus</i> (Butler, 1883)	*
<i>C. danae eupompe</i> (Klug, 1829)	*
<i>C. evarne</i> (Klug, 1829)	*
<i>C. vesta princeps</i> (Talbot 1939)	*
<i>C. regina</i> (Trimen, 1863)	*^
Genus <i>Eronia</i> Hübner, 1823	
<i>E. cleodora</i> Hübner, 1823	
Genus <i>Afrodryas</i> Stoneham, 1957	
<i>A. leda</i> (Boisduval, 1847)	
Tribe Pierini Swainson, 1820	
Genus <i>Appias</i> Hübner, 1819	
<i>A. sabina sabina</i> (Felder & Felder, 1865)	
<i>A. phaola intermedia</i> Dufrane, 1848	
<i>A. epaphia epaphia</i> (Cramer, 1779)	
Genus <i>Belenois</i> Hübner, 1819	
<i>B. creona severina</i> (Stoll, 1781)	
<i>B. aurota</i> (Fabricius, 1793)	

<i>B. crawshayi</i> Butler, 1894	
<i>B. calypso marlieri</i> Berger, 1981	
<i>B. thysa meldolae</i> Butler, 1872	
<i>B. theora laeta</i> (Weymer, 1903)	
<i>B. solilucis loveni</i> (Aurivillius, 1921)	
<i>B. sudanensis katalensis</i> Berger, 1981	*
<i>B. subeida subeida</i> (Felder & Felder, 1865)	*
Genus <i>Dixeia</i> Talbot, 1932	
<i>D. pigea</i> (Boisduval, 1836)	
Genus <i>Mylothris</i> Hübner, 1819	
<i>M. chloris clarissa</i> (Butler, 1888)	
<i>M. rhodope</i> (Fabricius, 1775)	
<i>M. continua continua</i> Talbot, 1944	
<i>M. agathina agathina</i> Cramer, 1779	*
Total for the subfamily Pierinae	36
Total for the family Pieridae	45
FAMILY RIODINIDAE Grote, 1895	
Subfamily Nemeobiinae Bates, 1868	
Tribe Abisarini Stichel, 1928	
Genus <i>Afriodinia</i> d'Abrera, 2009	
<i>A. neavei kivuensis</i> (Riley, 1932)	
Total for the family Riodinidae	1
FAMILY LYCAENIDAE Leach, 1815	
Subfamily Poritiinae Doherty, 1886	
Tribe Pentilini Aurivillius, 1914	
Genus <i>Ptelina</i> Clench, 1965	
<i>P. carnuta kamitugensis</i> (Dufrane, 1946)	
Genus <i>Pentila</i> Westwood, 1851	
<i>P. pauli clarensis</i> Neave, 1903	
<i>P. umangiana connectens</i> Hulstaert, 1924	
<i>P. inconspicua</i> Druce, 1910	
<i>P. tachyroides tachyroides</i> Dewitz, 1879	
<i>P. cloetensi catauga</i> Rebel, 1914	
<i>P. hiendlmayri</i> (Dewitz, 1887)	
<i>P. hybrida</i> Libert & Collins, 2019	*
Genus <i>Telipna</i> Aurivillius, 1895	
<i>T. aurivillii jefferyi</i> Jackson, 1969	
<i>T. sanguinea depuncta</i> Talbot, 1937	
<i>T. consanguinea consanguinea</i> Rebel, 1914	
<i>T. nyanza katangae</i> Stempffer, 1961	
<i>T. acraea nigrita</i> Jackson, 1969	*
<i>T. citrimaculata neavei</i> Bethune-Baker, 1926	*

<i>T. erica</i> Suffert, 1904	*
Genus <i>Ornipholidotos</i> Bethune-Baker, 1914	
<i>O. overlaeti intermedia</i> Libert, 2005	
<i>O. jacksoni jacksoni</i> Stempffer, 1961	
<i>O. ugandae ugandae</i> Stempffer, 1947	*
<i>O. ntebi</i> Bethune-Baker, 1906	*
<i>O. paradoxa orientis</i> Libert, 2005	*
Tribe Liptenini Röber, 1892	
Genus <i>Liptena</i> Westwood, 1851	
<i>L. ugandana</i> Stempffer, Bennett & May	
<i>L. xanthostola xantha</i> (Grose-Smith, 1901)	
<i>L. kamitugensis</i> Dufrane, 1945	
<i>L. flavicans praeusta</i> Schultze, 1917	*
<i>L. introspectionem</i> Szabolcs, 2021	*
<i>L. hapale</i> Talbot, 1935	*
Genus <i>Tetrarhanis</i> Karsch, 1893	
<i>T. ilala ilala</i> (Riley, 1929)	^
<i>T. ilma ugandae</i> (Stempffer, 1964)	
Genus <i>Falcuna</i> Stempffer & Bennett, 1963	
<i>F. margarita</i> (Suffert, 1904)	
<i>F. orientalis bwamba</i> Stempffer & Bennett, 1963	
<i>F. iturina</i> Stempffer & Bennett, 1963	*
<i>F. kasai</i> Stempffer & Bennett, 1963	*
Genus <i>Larinopoda</i> Butler, 1871	
<i>L. tera</i> (Hewitson, 1873)	
<i>L. lagyra lagyra</i> (Hewitson 1866)	
Genus <i>Micropentila</i> Aurivillius, 1895	
<i>M. fontainei</i> Stempffer & Bennett, 1965	*
Genus <i>Citrinophila</i> Kirby, 1887	
<i>C. erastus pallida</i> Hawker-Smith, 1933	
<i>C. tenera</i> (Kirby, 1887)	*
Genus <i>Toxochitona</i> Stempffer, 1956	
<i>T. sankuru</i> Stempffer, 1961	
Genus <i>Mimacraea</i> Butler, 1872	
<i>M. landbecki</i> Druce, 1910	
<i>M. krausei krausei</i> Dewitz, 1889	
<i>M. fulvaria fulvaria</i> Aurivillius, 1895	*
<i>M. telloides</i> Schultz, 1923	*
Genus <i>Mimeresia</i> Stempffer, 1961	
<i>M. drucei ugandae</i> (Stempffer, 1954)	
<i>M. moreelsi purpurea</i> (Hawker-Smith, 1933)	
Tribe Epitolini Jackson, 1962	

Genus <i>Iridana</i> Aurivillius, 1920	
<i>I. bwamba</i> Stempffer, 1964	*^
<i>I. obscura</i> Stempffer, 1964	*^
<i>I. katera</i> Stempffer, 1964	*
Genus <i>Epitola</i> Westwood, 1851	
<i>E. uranoides uranoides</i> Libert, 1999	
<i>E. urania</i> Kirby, 1887	
Genus <i>Cerautola</i> Libert, 1999	
<i>C. mittoni</i> (Jackson, 1964)	
<i>C. subargentea subargentea</i> (Jackson, 1965)	
<i>C. crowleyi holochroma</i> (Berger, 1981)	*
Genus <i>Geritola</i> Libert, 1999	
<i>G. cynaea</i> (Jackson, 1964)	
<i>G. liana</i> (Roche, 1954)	*
<i>G. dubia dubia</i> (Jackson, 1964)	*
<i>G. zelica viridis</i> (Libert, 2020)	*
Genus <i>Stempfferia</i> Jackson	
<i>S. cercenoides</i> (Holland, 1890)	
<i>S. tumentia</i> (Druce, 1910)	
<i>S. badura contrasta</i> Libert, 1999	*
<i>S. cercene</i> (Hewitson, 1873)	*
<i>S. ciconia mongiro</i> (Jackson, 1968)	*
<i>S. congoana orientalis</i> (Libert, 2020)	*
<i>S. insulana</i> (Aurivillius, 1923)	*
<i>S. sylviae</i> Libert, 1999	*
Genus <i>Cephetola</i> Libert, 1999	
<i>C. bwamba</i> (Jackson, 1964)	
<i>C. pinodes budduana</i> (Talbot, 1937)	
<i>C. entebbeana</i> (Bethune-Baker, 1926)	*
<i>C. cephenoides</i> Libert, 2020	*
<i>C. maculata</i> (Hawker-Smith, 1926)	*
<i>C. viridana</i> (Joicey and Talbot, 1921)	*
<i>C. orientalis</i> (Roche, 1954)	*
<i>C. kamengensis subgriseata</i> (Jackson, 1964)	*
<i>C. mpangensis mpangensis</i> (Jackson, 1962)	*
<i>C. dolorosa</i> Roche, 1954	*
Genus <i>Epitolina</i> Aurivillius, 1895	
<i>E. dispar</i> (Kirby, 1887)	
<i>E. melissa</i> (Druce, 1888)	*
Genus <i>Phytala</i> Westwood, 1851	
<i>P. elais ugandae</i> Jackson, 1964	
Genus <i>Hewitsonia</i> Kirby, 1871	

<i>H. ugandae ugandae</i> Jackson, 1962	
<i>H. inexpectata</i> Bouyer, 1997	
<i>H. intermedia</i> Jackson, 1962	
<i>H. kuehnei</i> Collins & Larsen, 2008	*^
Genus Powellana Bethune-Baker, 1908	
<i>P. cottoni</i> Bethune-Baker, 1908	*
Total for the subfamily Poritiinae	82
Subfamily Miletinae Reuter, 1896	
Tribe Lachnocnemini Clench, 1955	
Genus Lachnocnema Trimen, 1887	
<i>L. exiguus</i> Holland, 1890	
Tribe Spalgini Toxopeus, 1929	
Genus Spalgis Moore, 1879	
<i>S. jacksoni jacksoni</i> Stempffer, 1967	
Tribe Miletini Reuter, 1896	
Genus Megalopalpus Röber, 1886	
<i>M. simplex</i> Röber, 1886	
<i>M. metaleucus</i> Karsch, 1893	
<i>M. zymna</i> (Westwood, 1851)	
Total for the subfamily Miletinae	5
Subfamily Aphnaeinae Distant, 1884	
Genus Aphnaeus Hübner, 1819	
<i>A. orcas</i> (Drury, 1782)	
Genus Pseudaletis Druce, 1888	
<i>P. antimachus</i> (Staudinger, 1887)	*
Genus Cigaritis Donzel, 1848	
<i>C. apelles kiellandi</i> (Bouyer, 2011)	*^
<i>C. crustaria</i> (Holland, 1890)	*
Genus Zeritis Boisduval 1836	
<i>Zeritis neriene</i> Boisduval, 1836	*^
Total for the subfamily Aphnaeinae	5
Subfamily Polyommatainae Swainson, 1827	
Tribe Lycaenesthini Toxopeus, 1929	
Genus Anthene Doubleday, 1847	
<i>A. larydas</i> (Cramer, 1780)	
<i>A. crawshayi crawshayi</i> (Butler, 1899)	
<i>A. ligures ligures</i> (Hewitson, 1874)	
<i>A. sylvanus</i> (Drury, 1773)	
<i>A. rubricinctus rubricinctus</i> (Holland, 1891)	
<i>A. ituria</i> (Bethune-Baker, 1910)	
<i>A. princeps</i> (Butler, 1876)	
<i>A. liodes liodes</i> (Hewitson, 1874)	

<i>A. lunulata grosei</i> (Aurivillius, 1899)	*
<i>A. akoae albidior</i> Libert, 2010	*
<i>A. irumu</i> (Stempffer, 1948)	*^
<i>A. eliasi</i> Congdon, Kielland & Collins, 1998	*
Genus <i>Neurellipes</i> Bethune-Baker, 1910	
<i>N. fulvimacula</i> (Mabille, 1890)	
<i>N. kampala triangularis</i> Libert, 2010	
<i>N. scintillula scintillula</i> (Holland, 1891)	
<i>N. makala</i> (Bethune-Baker, 1910)	*
<i>N. zenkeri zenkeri</i> (Karsch, 1895)	*
<i>N. lachares toroensis</i> (Stempffer, 1947)	*
<i>N. onias</i> (Hulstaert, 1924)	*
Genus <i>Triclema</i> Karsch, 1893	
<i>T. staudingeri</i> (Grose-Smith & Kirby, 1894)	
<i>T. phoenicis</i> Karsch, 1893	
<i>T. inconspicua latefascia</i> Libert, 2010	
<i>T. obsoleta</i> (Stempffer, 1947)	*
<i>T. rufoplagata ituriensis</i> Joicey & Talbot, 1921	*
<i>T. lamias katerae</i> d'Abrera, 1980	*
Tribe Polyommata Swainson, 1827	
Genus <i>Cupidopsis</i> Karsch, 1895	
<i>C. cissus cissus</i> (Godart, 1824)	
Genus <i>Pseudonacaduba</i> Stempffer, 1943	
<i>P. aethiops</i> (Mabille, 1877)	
<i>P. sichela sichela</i> Wallengren, 1857)	
Genus <i>Uranothauma</i> Butler, 1895	
<i>U. falkensteinii</i> (Dewitz, 1879)	
<i>U. cyara tenuimarginata</i> (Grünberg, 1908)	
<i>U. heritsia intermedia</i> (Tite, 1958)	*
Genus <i>Cacyreus</i> Butler, 1897	
<i>C. lingeus</i> (Stoll, 1782)	
<i>C. audeoudi</i> Stempffer, 1936	
Genus <i>Leptotes</i> Scudder, 1876	
<i>L. pirithous pirithous</i> (Linnaeus, 1767)	
<i>L. marginalis</i> (Stempffer, 1942)	
Genus <i>Tuxentius</i> Larsen, 1982	
<i>T. margaritaceus</i> (Sharpe, 1892)	
Genus <i>Zizeeria</i> Chapman, 1910	
<i>Z. knysna knysna</i> (Trimen, 1862)	
Genus <i>Actizera</i> Chapman, 1910	
<i>A. lucida</i> (Trimen, 1883)	
Genus <i>Zizula</i> Chapman, 1910	

<i>Z. hylax</i> (Fabricius, 1775)	
Genus <i>Azanus</i> Moore, 1881	
<i>A. isis</i> Drury, 1773	
<i>A. mirza</i> (Plötz, 1880)	
<i>A. moriqua</i> (Wallengren, 1857)	^
<i>A. natalensis</i> (Trimen, 1887)	
Genus <i>Eicochrysops</i> Bethune-Baker, 1924	
<i>E. hippocrates</i> (Fabricius, 1793)	
Genus <i>Euchrysops</i> Butler, 1900	
<i>E. osiris</i> (Hopffer, 1855)	
<i>E. barkeri</i> (Trimen, 1893)	
Genus <i>Thermoniphas</i> Karsch, 1895	
<i>T. alberici</i> (Dufrane, 1945)	
<i>T. distincta</i> (Talbot, 1935)	^
<i>T. fontainei</i> Stempffer, 1956	^
<i>T. togara togara</i> Plötz, 1880	*
Genus <i>Oboronia</i> Karsch, 1893	
<i>O. punctatus</i> (Dewitz, 1879)	
<i>O. pseudopunctatus</i> (Strand, 1912)	
<i>O. guessfeldtii</i> (Dewitz, 1879)	
<i>O. ornata vestalis</i> (Aurivillius, 1895)	*^
<i>O. albicosta</i> (Gaede, 1916)	*
Tribe Hypolycaenini Swinhoe, 1910	
Genus <i>Hypolycaena</i> Felder, 1862	
<i>H. antifaunus latimacula</i> (Joicey & Talbot, 1921)	
<i>H. philippus philippus</i> (Fabricius, 1793)	
<i>H. nigra</i> Bethune-Baker, 1914	
<i>H. hatita ugandae</i> Sharpe, 1904	
Total for the subfamily Polyommatainae	59
Subfamily Theclinae Swainson, 1830	
Tribe Amblypodini Doherty, 1886	
Genus <i>Myrina</i> Fabricius, 1807	
<i>M. sharpei sharpei</i> Bethune-Baker, 1906	
Tribe Iolaini Riley, 1958	
Genus <i>Iolais</i> Hübner, 1819	
<i>I. bellina exquisita</i> (Riley, 1928)	
<i>I. pollux albocaerulea</i> (Riley, 1929)	
<i>I. mongiro</i> (Stempffer, 1969)	*^
<i>I. creta</i> (Hewitson, 1878)	*
<i>I. neavei katera</i> (Talbot, 1937)	*
Tribe Hypolycaenini Swinhoe, 1910	
Tribe Oxyliadini Eliot, 1973	

Genus <i>Oxylides</i> Hübner, 1819	
<i>O. albata</i> (Aurivillius, 1895)	^
<i>O. feminina feminina</i> (Sharpe, 1904)	
Tribe Deudorini Doherty, 1886	
Genus <i>Deudorix</i> Hewitson, 1863	
<i>D. antalus</i> (Hopffer, 1855)	
<i>D. odana odana</i> Druce, 1887	*
Genus <i>Hypomyrina</i> Druce, 1891	
<i>H. nomenia</i> (Hewitson, 1874)	*
<i>H. fournerae</i> Gabriel, 1939	*
Genus <i>Paradeudorix</i> Libert, 2004	
<i>P. canescens</i> (Joicey & Talbot, 1921)	*
Genus <i>Pilodeudorix</i> Libert, 2004	
<i>P. congoana orientalis</i> (Stempffer, 1957)	*
<i>P. anetia</i> (Hulstaert, 1924)	*
<i>P. mera mera</i> (Hewitson, 1873)	*
<i>P. otraeda genuba</i> (Hewitson, 1875)	*
<i>P. dimetris indentata</i> Libert, 2004	*^
<i>P. bwamba</i> (Stempffer, 1962)	*^
<i>P. violetta</i> (Aurivillius, 1897)	*^
<i>P. cameronia katanga</i> (Clench, 1965)	*
Total for the subfamily Theclinae	21
Total for the family Lycaenidae	172
Species total for Semuliki National Park	631

Taxon	Summary of recorded species
FAMILY PAPILIONIDAE Latreille, 1802	19
FAMILY HESPERIIDAE Latreille, 1809	109
FAMILY NYMPHALIDAE Rafinesque, 1815	285
FAMILY PIEIRIDAE Swainson, 1820	45
FAMILY RIODINIDAE Grote, 1895	1
FAMILY LYCAENIDAE Leach, 1815	172
Taxon total for Semuliki National Park	631

- * new record (additional to Forbes, 2018)
- ^ not recorded at DRC Ituri or Semliki Valley sampling sites (Ducarme, 2024)
- + *B.alboplaga* and *B. xenoides* treated as a single species (Aduse-Poku *et al.*, 2017)

References and Notes (by Genus)

Family Papilionidae

Notes:

Species are grouped under subgenera and species groups within the website, the phylogeny used is Condamine, (2023).

References:

Condamine, F.L., Allio, R., Reboud, E.I. *et al.*, (2023). A comprehensive phylogeny and revised taxonomy illuminate the origin and diversification of the global radiation of *Papilio* (Lepidoptera: Papilionidae). *Molecular phylogenetics and evolution* 183.

Family HesperIIDae

Notes:

A number of new records have been discovered from the unpublished Larsen HesperIIDae document.

Species within this family have been grouped into tribes using referenced work by Zhang *et al.*, (2022) and Huang *et al.*, (2024). There are some differences in their systematic conclusions and in this species list the latest reference work will take precedence.

References:

Huang, Z., Chiba, H., Hu, Y., *et al.*, (2024). Molecular phylogeny of HesperIIDae (Lepidoptera) with an emphasis on Asian and African genera. *Molecular Phylogenetics and Evolution*, 198.

Larsen, T.B., (2015). A revision of the Family HesperIIDae (unpublished document), ABRI, Nairobi, Kenya.

Zhang, J., Cong, Q., Shen, J., & Grishin, N.V. (2022). Taxonomic changes suggested by the genomic analysis of HesperIIDae (Lepidoptera). *Insecta Mundi*, 921.

Family Nymphalidae

Notes:

The genus *Acraea* taxonomic classification uses that of Beranud and Pierre (2018).

The genus *Neptis* has followed the latest taxonomic revisions by Richardson (2019, 2020)

References:

Bernaudo, D., Lequeux, J-P. & Ziraye, M., (2019). *Acraeinae* of Uganda.

Richardson, I., (2019). Revision of the genus *Neptis* Fabricius, 1807 (Lepidoptera, Nymphalidae) in the Afrotropical Region: Currently described taxa. *Metamorphosis* 30, pp 69-221.

Richardson, I., (2020). Revision of the genus *Neptis* Fabricius, 1807 (Lepidoptera: Nymphalidae) in the Afrotropical Region, Part 2: Two new species in the Agatha group. *Metamorphosis* 31, pp 84-93.

Acknowledgements

Mark Williams individual genus documents (Afrotropical Lepidopterists' Society) were used for species distribution data. Special thanks are also due to Robert Ducarme for our discussions on shared subspecies and differences between our records from Semuliki and his recording locations at Semuliki Valley and the Ituri Forest of the DRC.

I am also very grateful to the following Afrotropical butterfly taxonomic specialists for their discussions and contributions in providing accurate identification confirmations when requested:

Regional species and subspecies collaboration - Robert Ducarme and Mark Williams

Subfamily Acraeinae - Dominique Bernaud

Genus *Euphaedra* - Gilles Faravel and Milan Zubrik

Genus *Neptis* - Ian Richardson

Family HesperIIDae - Michel Libert

Family Pieridae - Haydon Warren-Gash

Family Lycaenidae - Michel Libert and Harald Selb